

409.

ON THE CONDITIONS FOR THE EXISTENCE OF THREE EQUAL ROOTS, OR OF TWO PAIRS OF EQUAL ROOTS, OF A BINARY QUARTIC OR QUINTIC.

[From the *Philosophical Transactions of the Royal Society of London*, vol. CLVIII. (for the year 1868), pp. 577—588. Received November 26, 1867,—Read January 9, 1868.]

[It is remarked, *Proc. R. Soc.* vol. XVII. p. 314, that the above title is a misnomer: I had in fact in regard to the quintic considered *not* the twofold relations belonging to the root-systems 311 and 221 respectively, but the threefold relations belonging to the root-systems 41 and 32 respectively. The proper title would have been "On the conditions for the existence of certain systems of equal roots of a binary quartic or quintic."]

In considering the conditions for the existence of given systems of equalities between the roots of an equation, we obtain some very interesting examples of the composition of relations. A relation is either onefold, expressed by a single equation $U=0$, or it is, say k -fold, expressed by a system of k or more equations. Of course, as regards onefold relations, the theory of the composition is well known: the relation $UV=0$ is a relation compounded of the relations $U=0$, $V=0$; that is, it is a relation satisfied if, and not satisfied unless, one or the other of the two component relations is satisfied. The like notion of composition applies to relations in general; viz., the compound relation is a relation satisfied if, and not satisfied unless, one or the other of the two component relations is satisfied. I purposely refrain at present from any further discussion of the theory of composition. I say that the conditions for the existence of given systems of equalities between the roots of an equation furnish instances of such composition; in fact, if we express that the function $(*\chi x, y)^n$, and its first-derived function in regard to x , or, what is the same thing, that the first-derived functions in regard to x, y respectively, have a common quadric factor, we obtain between the coefficients a certain twofold relation, which implies either that the equation $(*\chi x, y)^n=0$ has three equal roots, or else that it has two pairs of equal roots; that is, the relation in question is satisfied if, and it is not satisfied

unless, there is satisfied either the relation for the existence of three equal roots, or else the relation for the existence of two pairs of equal roots; or the relation for the existence of the quadric factor is compounded of the last-mentioned two relations. The relation for the quadric factor, for any value whatever of n , is at once seen to be expressible by means of an oblong matrix, giving rise to a series of determinants which are each to be put = 0; the relation for three equal roots and that for two pairs of equal roots, in the particular cases $n=4$ and $n=5$, are given in my "Memoir on the Conditions for the existence of given Systems of Equalities between the roots of an Equation," *Phil. Trans.* vol. CXLVII. (1857), pp. 727—731, [150]; and I propose in the present Memoir to exhibit, for the cases in question $n=4$ and $n=5$, the connexion between the compound relation for the quadric factor with the component relations for the three equal roots and for the two pairs of equal roots respectively.

Article Nos. 1 to 8, *the Quartic.*

1. For the quartic function

$$(a, b, c, d, e\zeta(x, y)^4,$$

the condition for three equal roots, or, say, for a root system 31, is that the quadric-variant and the cubinvariant each of them vanish, viz. we must have

$$\begin{aligned} I &= ae - 4bd + 3c^2 = 0, \\ J &= ace - ad^2 - b^2e + 2bcd - c^3 = 0. \end{aligned}$$

2. The condition for two pairs of equal roots, or for a root system 22, is that the cubicovariant vanishes identically, viz. representing this by

$$(A, B, 5C, 10D, 5E, F, G^5\zeta(x, y)^6 = 0,$$

we must have

$$\begin{aligned} A &= a^2d - 3abc + 2b^3 = 0, \\ B &= a^2e + 2abd - 9ac^2 + 6b^2c = 0, \\ C &= abe - 3acd + 2b^2d = 0, \\ D &= -ad^2 + b^2e = 0, \\ E &= -ade + 3bce - 2bd^2 = 0, \\ F &= -ae^2 - 2bde + 9c^2e - 6cd^2 = 0, \\ G &= -be^2 + 3cde - 2d^3 = 0. \end{aligned}$$

3. But the condition for the common quadric factor is

$$\begin{vmatrix} a, & 3b, & 3c, & d \\ b, & 3c, & 3d, & e \\ a, & 3b, & 3c, & d \\ b, & 3c, & 3d, & e \end{vmatrix} = 0,$$

and the determinants formed out of this matrix must therefore vanish for $(I, J)=0$, and also for $(A, B, C, D, E, F, G)=0$, that is, the determinants in question must be syzygetically related to the functions (I, J) , and also to the functions (A, B, C, D, E, F, G) .

4. The values of the determinants are

1234 = 3 ×	1235 = 3 ×	1245 =	1345 = 3 ×	2345 = 3 ×
$a^2ce + 1$	$a^2de - 1$	$a^2e^2 - 1$	$abe^2 - 1$	$ace^2 + 1$
$a^2d^2 - 3$	$abce + 4$	$abde + 2$	$acde + 4$	$ad^2e - 1$
$ab^2e - 1$	$abd^2 + 1$	$ac^2e + 9$	$ad^3 - 3$	$b^2e^2 - 3$
$abcd + 14$	$ac^2d - 3$	$acd^2 - 9$	$b^3de + 1$	$bcde + 14$
$ac^3 - 9$	$b^3e - 3$	$b^2ce - 9$	$bc^2e - 3$	$bd^3 - 8$
$b^3d - 8$	$b^2cd + 2$	$b^2d^2 + 8$	$bcd^2 + 2$	$c^3e - 9$
$b^2c^2 + 6$				$c^2d^2 + 6$

5. The syzygetic relation with (I, J) is given by means of the identical equation

$$\begin{vmatrix} y^4, & -4xy^3, & 6x^2y^2, & -4x^3y, & x^4 \\ & a, & 3b, & 3c, & d \\ & b, & 3c, & 3d, & e \\ a, & 3b, & 3c, & d, & \\ b, & 3c, & 3d, & e, & \end{vmatrix} = -6I \cdot \tilde{H}U + 9J \cdot U,$$

or, as this may be written,

$$(1234, 1235, 1245, 1345, 2345)\tilde{\chi}(x, y)^4 = -6I \cdot \tilde{H}U + 9J \cdot U,$$

where $\tilde{H}U$ is the Hessian of U ,

$$= \begin{vmatrix} ac + 1 & ad + 2 & ae + 1 & be + 2 & ce + 1 \\ b^2 - 1 & bc - 2 & bd + 2 & cd - 2 & d^2 - 1 \\ & & c^2 - 3 & & \end{vmatrix} \tilde{\chi}(x, y)^4.$$

6. That is, we have

$$\begin{aligned} 1234 &= (ac - b^2, a\tilde{\chi} - 6I, 9J), \\ 4 \cdot 1235 &= (2ad - 2bc, 4b\tilde{\chi} - 6I, 9J), \\ 6 \cdot 1245 &= (ae + 2bd - 3c^2, 6c\tilde{\chi} - 6I, 9J), \\ 4 \cdot 1345 &= (2be - 2cd, 4d\tilde{\chi} - 6I, 9J), \\ 2345 &= (ce - d^2, e\tilde{\chi} - 6I, 9J). \end{aligned}$$

7. The determinants thus vanish if $(I, J) = 0$, that is, for the root system 31; they will also vanish without this being so, if only

$$\left(\frac{3J}{2I}\right) \frac{ac - b^2}{a} = \frac{ad - bc}{2b} = \frac{ae + 2bd - 3c^2}{6c} = \frac{be - cd}{2d} = \frac{ce - d^2}{e};$$

and we may omit the first member $\left(\frac{3J}{2I}\right)$, since if the remaining terms are equal

to each other they will also be $= \frac{3J}{2I}$. The equations may then be written

$$\begin{vmatrix} ac - b^2, & ad - bc, & ae + 2bd - 3c^2, & be - cd, & ce - d^2 \\ a, & 2b, & 6c, & 2d, & e \end{vmatrix} = 0,$$

and the ten equations of this system reduce themselves (as it is very easy to show) to the seven equations

$$(A, B, C, D, E, F, G) = 0,$$

which, as above mentioned, are the conditions for the root system 22.

8. It may be added that we have

	A	B	C	D	E	F	G
$\frac{1}{3} \cdot 1234 =$		c	$-4b$	$+3a$			
$\frac{1}{3} \cdot 1235 =$			c	$-3b$	$+a$		
$0 =$		d	$-3c$		$+a$		
$1245 =$		$-e$	$+4d$	$-3c$			
$0 =$		$-e$		$+6c$		$-a$	
$0 =$			$-d$	$+3c$	$-b$		
$\frac{1}{3} \cdot 1345 =$			$-e$	$+3d$	$-c$		
$0 =$			$-e$		$+3c$	$-b$	
$\frac{1}{3} \cdot 2345 =$				$-3e$	$+4d$	$-c$	

where it is to be noticed that the four equations having the left-hand side $= 0$, give $B : C : D : E : F$ proportional to the determinants of the matrix

$$\begin{vmatrix} d, & -3c, & ., & a \\ -e, & ., & 6c, & ., & -a \\ & -d, & 3c, & -b \\ & -e, & ., & +3c, & -b \end{vmatrix};$$

the determinants in question contain each the factor c , and omitting this factor, the system shows that B, C, D, E, F are proportional to their before-mentioned actual values.

Article Nos. 9 to 15, *the Quintic*.

9. For the quintic function

$$(a, b, c, d, e, f)(x, y)^5,$$

the condition of a root system 41 is that the covariant, $[B =]$ No. 14, shall vanish, viz. we must have

$$\begin{aligned} A &= 2(ae - 4bd + 3c^2) = 0, \\ B &= af - 3be + 2cd = 0, \\ C &= 2(bf - 4ce + 3d^2) = 0. \end{aligned}$$

10. The condition of a root system 32 is that the following covariant, viz.

$$[3A^2B - 25C^2, =] 3(\text{No. 13})^2(\text{No. 14}) - 25(\text{No. 15})^2,$$

shall vanish, where

[A =] No. 13 = $(a, b, c, d, e, f \chi(x, y))^5$, the quintic itself.

$$[B =] \text{ No. 14} = \begin{pmatrix} ae & bf \\ bd - 4 & ce - 4 \\ c^2 + 3 & cd + 2 \\ & d^2 + 3 \end{pmatrix} \chi(x, y)^2.$$

$$[C =] \text{ No. 15} = \begin{pmatrix} ac & ad & ae & bf \\ b^2 - & bc - 3 & bd + 3 & ce + 3 \\ & c^2 - 6 & cd - 6 & d^2 - 6 \end{pmatrix} \chi(x, y)^6.$$

11. The developed expression of the foregoing function is as follows :

\mathfrak{A}	\mathfrak{B}	\mathfrak{C}	\mathfrak{D}	\mathfrak{E}	\mathfrak{F}	\mathfrak{G}	\mathfrak{H}	\mathfrak{I}	\mathfrak{J}	\mathfrak{K}	\mathfrak{L}	\mathfrak{M}	
$a^2e + 3$	$a^2f + 3$	$a^2bf + 33$	$a^2ef + 10$	$a^2df - 90$	$a^2ef - 114$	$a^2f^2 - 114$	$abf^2 - 114$	$abf^2 - 114$	$acf^2 - 90$	$acf^2 - 90$	$adf^2 + 10$	$adf^2 + 10$	$af^3 + 3$
$a^2bd - 12$	$a^2be + 21$	$a^2ce - 102$	$a^2de - 390$	$a^2e^2 - 195$	$abd^2 - 264$	$abef - 264$	$acef - 264$	$acef - 264$	$adef + 360$	$adef + 360$	$ae^2f + 155$	$ae^2f + 155$	$bf^3 + 3$
$a^2c^2 - 16$	$a^2cd - 144$	$a^2d^2 - 216$	$ab^2f + 155$	$abcf + 360$	$abe^2 - 990$	$acd^2 + 990$	$acd^2 + 537$	$acd^2 + 537$	$ae^3 + 225$	$ae^3 + 225$	$bcf^2 - 390$	$bcf^2 - 390$	$cef^3 - 12$
$ab^2c + 50$	$ab^2d + 30$	$ab^2e + 135$	$abce + 100$	$abde - 1500$	$ae^2f + 468$	$ace^2 - 245$	$ace^2 - 245$	$ace^2 - 245$	$b^2f^2 - 195$	$b^2f^2 - 195$	$bdef + 100$	$bdef + 100$	$d^2f^2 - 16$
$b^4 - 25$	$abc^2 + 240$	$abcd + 120$	$abd^2 - 600$	$ac^2e + 900$	$acde + 1320$	$ad^2e + 1740$	$ad^2e + 1740$	$ad^2e + 1740$	$bce^2f - 1500$	$bce^2f - 1500$	$be^3 + 125$	$be^3 + 125$	$e^4 - 25$
	$b^2c - 150$	$ac^3 + 480$	$ac^2d + 1600$	$acd^2 + 1800$	$ad^3 + 1080$	$b^2df - 245$	$bcd^2 + 1320$	$bcd^2 + 1320$	$b^2e^2 - 1700$	$b^2e^2 - 1700$	$c^2ef - 600$	$c^2ef - 600$	
		$b^3d - 150$	$b^2e + 125$	$b^3f + 225$	$b^2ef + 900$	$b^2e^2 - 1700$	$bce^2 - 2700$	$bce^2 - 2700$	$c^2df + 1800$	$c^2df + 1800$	$cd^2f + 1600$	$cd^2f + 1600$	
		$b^2e^2 - 300$	$b^2cd - 1000$	$b^2d^2 - 1500$	$b^2de - 2700$	$bc^2f + 1740$	$bd^2e + 900$	$bd^2e + 900$	$c^2e^2 - 1500$	$c^2e^2 - 1500$	$cd^2e^2 - 1000$	$cd^2e^2 - 1000$	
					$bc^2e + 900$	$bcde - 2000$	$c^3f + 1080$	$c^3f + 1080$					
					$bcd^2 - 600$	$bd^3 + 600$	$c^2de - 600$	$c^2de - 600$					
						$c^2e + 600$							
						$c^2d^2 - 400$							

15. Assuming the existence of these relations, we have for the determination of the numerical coefficients in each relation a set of linear equations, which are shown by the following Tables, viz. referring to the Table headed $c\mathfrak{A}$, $b\mathfrak{B}$, $a\mathfrak{C}$, $a.1234$, [first of the seven tables *infra*] if the multipliers of the several terms respectively be A , B , C , X , then the Table denotes the system of linear equations

$$\begin{aligned} 0 A + 3 B + 33 C + 0 X &= 0, \\ 3 A + 0 B - 102 C - 16 X &= 0, \\ &\&c., \end{aligned}$$

that is, nine equations to be satisfied by the ratios of the coefficients A , B , C , X , and which are in fact satisfied by the values at the foot of the Table, viz.

$$A : B : C : X = +66 : -11 : +1 : +6.$$

There would be in all fourteen Tables, but as those for the second seven would be at once deducible by symmetry from the first seven, I have only written down the seven Tables; the solutions for the first and second Tables were obtained without difficulty, but that for the third Table was so laborious to calculate, and contains such extraordinarily high numbers, that I did not proceed with the calculation, and it is accordingly only the first, second, and third Tables which have at the foot of them respectively the solutions of the linear equations.

16. The results given by these three Tables are, of course,

$$\begin{aligned} 66 c\mathfrak{A} - 11 b\mathfrak{B} + 1 a\mathfrak{C} + 6 a.1234 &= 0, \\ 330 d\mathfrak{A} + 110 c\mathfrak{B} - 55 b\mathfrak{C} + 9 a\mathfrak{D} - 105 a.1235 &= 0, \\ + 266478575 e\mathfrak{A} & \\ - 617359490 d\mathfrak{B} & \\ + 144200810 c\mathfrak{C} & \\ + 9656911 b\mathfrak{D} & \\ + 9090785 a\mathfrak{E} & \\ - 721004050 c.1234 & \\ + 90914175 b.1235 & \\ - 160758675 a.1245 & \\ + 11559295 a.1236 &= 0. \end{aligned}$$

It is to be noticed that the nine coefficients of this last equation were obtained from, and that they actually satisfy, a system of fourteen linear equations; so that the correctness of the result is hereby verified.

17. The seven Tables are

First Table.

	cA	bB	aC	$a.1234$
a^3bf		+ 3	+ 33	
a^3ce	+ 3		- 102	- 16
a^3d^2			- 216	+ 36
a^2be^2		+ 21	+ 135	+ 16
a^2bcd	- 12	- 144	+ 120	- 152
a^2c^3	- 16		+ 480	+ 96
ab^3d		+ 30	- 150	+ 80
ab^2c^2	+ 50	+ 240	- 300	- 60
b^4c	- 25	- 150		
	+ 66	- 11	+ 1	+ 6

Second Table.

	dA	eB	bC	aD	$a.1235$	$b.1234$
a^3cf		+ 3		+ 10	- 4	
a^3de	+ 3			- 390	+ 24	
a^2b^2f			+ 33	+ 155	+ 4	
a^2bce		+ 21	- 102	+ 100	- 84	- 16
a^2bd^2	- 12		- 216	- 600	- 24	+ 36
a^2c^2d	- 16	- 144		+ 1600	+ 64	+ 16
ab^3e			+ 135	+ 125	+ 60	
ab^2cd	+ 50	+ 30	+ 120	- 1000	- 40	- 152
abc^3		+ 240	+ 480			+ 96
b^4d	- 25		- 150			+ 80
b^3c^2		- 150	- 300			- 60
	+ 330	+ 110	- 55	+ 9	- 105	0

Third Table.

	eA	dB	cC	bD	aE	$c.1234$	$b.1235$	$a.1245$	$a.1236$
a^3df		+ 3			- 90			- 6	+ 6
a^3e^2	+ 3				- 195			+ 16	
a^2bcf			+ 33	+ 10	+ 360		- 4	+ 6	- 22
a^2bde	- 12	+ 21		- 390	- 1500		+ 24	- 26	- 6
a^2c^2e	- 16		- 102		+ 900	- 16		- 96	+ 16
a^2cd^2		- 144	- 216		+ 1800	+ 36		+ 96	
ab^3f				+ 155	+ 225		+ 4		+ 16
ab^2ce	+ 50		+ 135	+ 100		+ 16	- 84	+ 90	- 10
ab^2d^2		+ 30		- 600	- 1500		- 24	- 80	
abc^2d		+ 240	+ 120	+ 1600		- 152	+ 64		
ac^4			+ 480			+ 96			
b^4e	- 25			+ 125			+ 60		
b^3cd		- 150	- 150	- 1000		+ 80	- 40		
b^3c^2			- 300			- 60			
	+266478575	-617359490	+144200810	+9656911	+9090785	-721004050	+90914175	-160758675	+11559295

Fourth Table.

	fA	eB	dC	cD	bE	aF	$d.1234$	$c.1235$	$b.1236$	$b.1245$	$a.1246$	$a.1345$
a^3ef	+ 3	+ 3				- 114					+ 4	
a^2bdf	- 12		+ 33		- 90	- 264			+ 6	- 6	- 4	- 24
a^2be^2		+ 21			- 195	- 990				+ 16	- 4	+ 64
a^2c^2f	- 16			+ 10		+ 468		- 4			- 24	+ 24
a^2cde		- 144	- 102	- 390		+ 1320	- 16	+ 24			+ 24	- 208
a^2d^3			- 216			+ 1080	+ 36					+ 144
ab^3cf	+ 50			+ 155	+ 360	+ 900		+ 4	- 22	+ 6	+ 24	
ab^2de		+ 30	+ 135		- 1500	- 2700	+ 16		- 6	- 26	- 20	- 40
abc^2e		+ 240		+ 100	+ 900	+ 900		- 84	+ 16	- 96		+ 60
$abcd^2$			+ 120	- 600	+ 1800	- 600	- 152	- 24		+ 96		- 40
ac^3d			+ 480	+ 1600			+ 96	+ 64				
b^4f	- 24				+ 225				+ 16			
b^3ce		- 150		+ 125				+ 60	- 10	+ 90		
b^3d^2			- 150		- 1500		+ 80			- 80		
b^2c^2d			- 300	- 1000			- 60	- 40				

Fifth Table.

	fB	eC	dD	cE	bF	aG	$e.1234$	$d.1235$	$c.1236$	$c.1245$	$b.1246$	$b.1345$	$a.1256$	$a.2345$	$a.1346$
a^3f^2	+ 3					- 19							+ 1		
a^2bef	+ 21	+ 33			- 114	- 608					+ 4		- 2		+ 16
a^2cdf	- 144		+ 10	- 90		+ 537	- 4	+ 6	- 6				- 16	+ 20	- 36
a^2ce^2		- 102	- 390	- 195		- 245	- 16		+ 16				+ 16	- 80	- 16
a^2d^3e		- 216			+ 1740	+ 1740	+ 36	+ 24					+ 16	+ 60	+ 36
ab^3df	+ 30		+ 155		- 264	- 245		+ 4			- 4	- 24	- 15	- 80	- 16
ab^2e^2		+ 135			- 990	- 1700	+ 16			- 4	+ 64			+ 240	
abc^2f	+ 240			+ 360	+ 468	+ 1740			- 22	+ 6	- 24	+ 24		+ 60	+ 36
$abcde$		+ 120	+ 100	- 1500	+ 1320	- 2000	- 152	- 84	- 6	- 26	+ 24	- 208		- 860	- 20
abd^3			- 600		+ 1080	+ 600		- 24				+ 144		+ 960	
ac^3e		+ 480		+ 900	+ 600	+ 600	+ 96		+ 16	- 96				+ 960	
ac^2d^2			+ 1600	+ 1800	- 400			+ 64		+ 96				- 320	
b^3cf	- 150			+ 225	+ 900				+ 16		+ 24				
b^2de		- 150	+ 125		- 2700		+ 80	+ 60			- 20	- 40			
b^2c^2e		- 300		+ 900			- 60		- 10	+ 90		+ 60			
b^2cd^2			- 1000	- 1500	- 600			- 40		- 80		- 40			

Sixth Table.

$f\mathbb{E}$	$e\mathbb{D}$	$d\mathbb{E}$	$e\mathbb{F}$	$b\mathbb{G}$	$a\mathbb{H}$	$f.1234$	$e.1235$	$d.1236$	$d.1245$	$e.1246$	$c.1345$	$b.1256$	$b.1346$	$a.2346$	$a.1356$
$a^2b^2c^2$	+ 33	-	- 114	-	19	- 114	- 16	- 4	+	4	+	+	-	24	+
a^2cef	- 102	-	-	-	-	- 264	+ 36	-	-	-	-	-	-	24	-
a^2d^2f	- 216	-	-	-	-	468	+ 36	+	+	+	+	+	+	24	-
a^2de^2	+ 135	-	-	-	698	900	16	+	+	+	+	+	+	64	+
ab^2ef	+ 120	-	-	-	537	1320	- 152	-	- 22	+	-	20	+	208	+
$abcd^2f$	+ 100	-	-	-	245	2700	- 84	-	4	+	+	80	-	40	+
abc^2e^2	+ 600	-	-	-	1740	900	- 24	-	-	+	+	60	+	60	-
abd^2e	+ 480	-	-	-	468	1080	96	+	+	- 24	+	24	+	144	+
ac^2f	+ 1600	+	+	+	1320	600	64	+	16	+	24	208	-	40	-
ac^2de	- 150	-	-	-	1080	-	60	+	96	+	144	80	-	40	-
acd^3	- 300	-	-	-	245	-	60	+	+	+	-	240	+	36	+
b^2df	+ 125	+	+	+	1700	900	- 60	-	24	+	40	60	+	20	+
b^2e^2f	- 1000	-	-	-	2000	600	- 40	-	- 20	-	-	860	-	960	-
b^2c^2de	- 1500	-	-	-	600	1500	- 1500	+	+	+	+	960	+	960	-
b^2d^3	- 1000	-	-	-	600	600	600	+	+	+	+	960	+	960	-
bc^2e	- 1000	-	-	-	600	600	600	+	+	+	+	960	+	960	-
bc^2d^2	- 1000	-	-	-	600	600	600	+	+	+	+	960	+	960	-

Seventh Table.

$f\mathbb{D}$	$e\mathbb{E}$	$d\mathbb{F}$	$e\mathbb{G}$	$b\mathbb{H}$	$a\mathbb{I}$	$f.1235$	$e.1245$	$d.1246$	$d.1345$	$e.1256$	$c.2345$	$b.2346$	$b.1356$	$a.2356$	$a.1456$
a^2ef^2	+ 10	-	-	-	90	- 24	+	+	+	+	+	+	+	6	+
a^2def	- 390	-	-	-	360	225	- 16	-	-	-	-	-	-	6	-
a^2e^3	- 195	-	-	-	225	900	+	+	+	+	+	+	+	16	+
ab^2f^2	+ 155	-	-	-	114	195	+	+	+	+	+	+	+	16	+
ab^2ef	+ 100	-	-	-	264	1500	- 22	-	24	+	16	24	+	26	+
abd^2f	+ 600	-	-	-	468	900	6	-	64	+	20	24	+	96	+
$abde^2$	- 1500	-	-	-	900	600	- 6	-	24	+	60	24	+	90	+
ac^2df	+ 1600	+	+	+	1800	1500	16	+	208	+	36	36	+	96	+
ac^2e^2	+ 900	+	+	+	1500	600	96	+	144	+	36	36	+	80	+
ac^2d^2e	+ 1800	+	+	+	1740	600	16	+	24	+	60	60	+	80	+
ad^4	+ 225	+	+	+	990	1320	16	+	24	+	80	64	+	80	+
b^2ef	+ 125	+	+	+	1320	2700	16	+	24	+	240	24	+	80	+
b^2cdf	+ 1000	-	-	-	245	2700	- 10	-	40	-	60	40	-	80	-
b^2ce^2	- 1500	-	-	-	1700	900	90	+	20	+	36	20	+	90	+
b^2d^2e	- 1500	-	-	-	1740	600	- 80	-	40	-	20	40	-	90	+
bc^2f	- 1000	+	+	+	1080	600	90	+	60	+	36	144	+	96	+
bc^2de	- 1000	+	+	+	600	600	- 10	-	40	-	20	40	-	80	+
bcd^3	- 1000	+	+	+	600	600	90	+	60	+	36	144	+	96	+
c^2e	- 1000	+	+	+	600	600	90	+	60	+	36	144	+	96	+
c^2d^2	- 1000	+	+	+	600	600	90	+	60	+	36	144	+	96	+

And the remaining seven Tables might of course be deduced from these by writing (f, e, d, c, b, a) instead of (a, b, c, d, e, f) , and making the corresponding alterations in the top line of each Table.

18. The equations $\mathfrak{A}=0, \mathfrak{B}=0, \dots, \mathfrak{M}=0$ consequently establish between the fifteen functions 1234, 1235, ... 3456 a system of fourteen equations, viz. the first and last three of these are

$$\begin{aligned} 1234 &= 0, \\ 1235 &= 0, \\ -160758675 \cdot 1245 \\ + 11559295 \cdot 1236 &= 0, \\ &\vdots \\ + 11559295 \cdot 1456 \\ -160758675 \cdot 2356 &= 0, \\ 2456 &= 0, \\ 3456 &= 0. \end{aligned}$$

To complete the proof that in virtue of the equations $\mathfrak{A}=0, \mathfrak{B}=0, \dots, \mathfrak{M}=0$ all the fifteen functions 1234, 1235, ... 3456 vanish, it is necessary to make use of the identical relations subsisting between these quantities 1234, &c.; thus we have

$$\begin{aligned} a \cdot 1345 + 4b \cdot 1245 + 6c \cdot 1235 + 4d \cdot 1234 &= 0, \\ b \cdot 1345 + 4c \cdot 1245 + 6d \cdot 1235 + 4e \cdot 1234 &= 0, \end{aligned}$$

which, in virtue of the above equations $1234=0$ and $1235=0$, become

$$\begin{aligned} a \cdot 1345 + 4b \cdot 1245 &= 0, \\ b \cdot 1345 + 4c \cdot 1245 &= 0, \end{aligned}$$

giving (unless indeed $ac-b^2=0$) $1245=0, 1345=0$; the equation $1245=0$ then reduces the third of the above equations to $1236=0$, and so on until it is shown that the fifteen quantities all vanish.