

# KEY FOR DETERMINATION OF POLISH LICHENS

by

Janusz NOWAK

The aim of this work is the development of a complete and modern key, much more detailed than that in "Manual of Polish lichens" of 1957, for determination of all species of lichens known to occur in Poland. Some taxa occurring in the adjacent territories of the German Democratic Republic, Southern Scandinavia and Czechoslovakia are included. Special attention is paid to the use of the current nomenclature of genera and species.

The manual will contain a total of approximately 2160 lichen species. Keys for the determination of species in the following genera have already been completed: *Absconditella*, *Acarospora*, *Acrocordia*, *Adeloclia*, *Agonimia*, *Alectoria*, *Allantoparmelia*, *Amygdalaria*, *Anaptychia*, *Anema*, *Anisomeridium*, *Anzina*, *Aphanoopsis*, *Arctonia*, *Arctoparmelia*, *Arthonia*, *Arthopyrenia*, *Arthorhaphis*, *Arthothelium*, *Arthrosporum*, *Aspicilia*, *Bacidia*, *Bactrospora*, *Baeomyces*, *Bagliettoa*, *Bellemerea*.

## References

1. Clauzade, G., Roux, C. 1985. Likenoj de okcidenta Eŭropo illustrata determinlibro. Bull. Soc. Bot. Centre-Ouest. num. special 7: 1-893.
2. Nowak, J., Tbolewski, Z. 1975. Manual of Polish lichens. (in Polish) pp. 1177. PWN, Warszawa.
3. Poelt, J., Vězda, A. 1977. Bestimmungsschlüssel europäischer Flechten. Ergänzungsheft I. Bibl. Lichenol. 9: 1-258.
4. Poelt, J., Vězda, A. 1981. Bestimmungsschlüssel europäischer Flechten. Ergänzungsheft II. Bibl. Lichenol. 16: 1-390.
5. Santesson, R. 1984. The Lichens of Sweden and Norway. pp. 333.
6. Wirth, V. Flechtenflora. UTB Ulmer. pp. 552.