

POPULATION BIOLOGY AND SEX RATIO OF DIOECIOUS PLANT SPECIES

by

Kazimierz ZARZYCKI & Urszula KORZENIAK

Studies on the sex ratio in populations of *Rumex acetosa* L. and *R. thysiflorus* Fingerh. were initiated many years ago (1-4). On the basis of both experimental and field observations, a simulation model of the changes in the sex ratio in *R. thysiflorus* has been prepared (in cooperation with J. Rychlewski, Jagiellonian University, and M. Dąmbrowski).

In addition, detailed study of the sex ratio in populations of *Anuncus silvester* Kost., *Antennaria dioica* (L.) Gaertn. and *Juniperus communis* L. are currently being carried out by U. Korzeniak.

References

1. Rychlewski, J., Zarzycki, K. 1975. Sex ratio in seeds of *Rumex acetosa* L. as a result of sparse or abundant pollination. *Acta Biol. Crac.*, Ser. Bot. 18: 101-114.
2. Rychlewski, J., Zarzycki, K. 1981. Sex ratio in seeds of *Rumex thysiflorus* Fingerh. from European and Canadian populations. *Acta Biol. Crac.*, Ser. Bot. 23: 97-105.
3. Rychlewski, J., Zarzycki, K. 1986. Genetical and ecological mechanisms regulating the sex ratio in populations of *Rumex thysiflorus* Fingerh. (*Polygonaceae*). *Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich* 87: 132-140.
4. Zarzycki, K., Rychlewski, J. 1972. Sex ratios in Polish natural populations and in seedling samples of *Rumex acetosa* L. and *R. thysiflorus* Fing. *Acta Biol. Crac.*, Ser. Bot. 15: 135-151.