















47. Mezey E., Key S., Vogelsang G. i wsp.: Transplanted bone marrow generates new neurons in human brains. *Proc. Natl. Acad. Sci. USA* 2003; 100 (3): 1364-1369.
48. Zhao L.R., Duan W.M., Reyes M. i wsp.: Human bone marrow stem cells exhibit neural phenotypes and ameliorate neurological deficits after grafting into the ischemic brain of rats. *Exp. Neurol.* 2002; 174 (1): 11-20.
49. Hess D.C., Hill W.D., Martin-Studdard A. i wsp.: Bone marrow as a source of endothelial cells and NeuN-expressing cells after stroke. *Stroke* 2002; 33 (5): 1362-1368.
50. Song S., Sanchez-Ramos J.: Preparation of neural progenitors from bone marrow and umbilical cord blood. *Methods Mol. Biol.* 2002; 198: 79-88.
51. Buzańska L., Machaj E.K., Zabłocka B. i wsp.: Human cord blood-derived cells attain neuronal and glial features *in vitro*. *J. Cell Sci.* 2002; 115 (Pt 10): 2131-2138.
52. Buzańska L., Stachowiak E., Stachowiak M. i wsp.: Neural stem cell line derived from human umbilical cord blood – morphological and functional properties. ESN abstrakt – w druku.
53. Buzańska L., Sun W., Salvi R.J. i wsp.: Changing electrophysiological properties in differentiating human cord blood-derived neural stem cells (HUCB-NSC). ESN abstrakt – w druku.
54. Andrews P.W.: Retinoic acid induces neuronal differentiation of a cloned human embryonal carcinoma cell line *in vitro*. *Dev. Biol.* 1984; 103 (2): 285-293.
55. Borlongan C.V., Tajima Y., Trojanowski J.Q. i wsp.: Transplantation of cryopreserved human embryonal carcinoma-derived neurons (NT2N cells) promotes functional recovery in ischemic rats. *Exp. Neurol.* 1998; 149 (2): 310-321.
56. Saporta S., Borlongan C.V., Sanberg PR: Neural transplantation of human neuroteratocarcinoma (hNT) neurons into ischemic rats. A quantitative dose-response analysis of cell survival and behavioral recovery. *Neuroscience* 1999; 91 (2): 519-525.
57. Kondziolka D., Wechsler L., Goldstein S. i wsp.: Transplantation of cultured human neuronal cells for patients with stroke. *Neurology* 2000; 55 (4): 565-569.
58. Baker K.A., Hong M., Sadi D., Mendez I.: Intrastratial and intranigral grafting of hNT neurons in the 6-OHDA rat model of Parkinson's disease. *Exp. Neurol.* 2000; 162 (2): 350-360.
59. Willing A.E., Janowski W., Igova T. i wsp.: A comparison of dopaminergic cells from the human Ntera2/D1 cell line transplanted into the hemiparkinsonian rat. *Exp. Neurol.* 2002; 175 (2): 447 INTR – 8. abstrakt.
60. Yoshimoto Y., Lin Q., Collier T.J. i wsp.: Astrocytes retrovirally transduced with BDNF elicit behavioral improvement in a rat model of Parkinson's disease. *Brain Res.* 1995; 691 (1-2): 25-36.
61. Frim D.M., Uhler T.A., Galpern W.R. i wsp.: Implanted fibroblasts genetically engineered to produce brain-derived neurotrophic factor prevent 1-methyl-4-phenylpyridinium toxicity to dopaminergic neurons in the rat. *Proc. Natl. Acad. Sci. USA* 1994; 91 (11): 5104-5108.
62. Levivier M., Przedborski S., Bencsics C., Kang U.J.: Intrastratial implantation of fibroblasts genetically engineered to produce brain-derived neurotrophic factor prevents degeneration of dopaminergic neurons in a rat model of Parkinson's disease. *J. Neurosci.* 1995; 15 (12): 7810-7820.
63. Arenas E., Trupp M., Akerud P., Ibanez C.F.: GDNF prevents degeneration and promotes the phenotype of brain noradrenergic neurons *in vivo*. *Neuron* 1995; 15 (6): 1465-1473.
64. Akerud P., Canals J.M., Snyder E.Y., Arenas E.: Neuroprotection through delivery of glial cell line-derived neurotrophic factor by neural stem cells in a mouse model of Parkinson's disease. *J. Neurosci.* 2001; 21 (20): 8108-8118.
65. Akerud P., Holm P.C., Castelo-Branco G. i wsp.: Persephin-overexpressing neural stem cells regulate the function of nigral dopaminergic neurons and prevent their degeneration in a model of Parkinson's disease. *Mol. Cell Neurosci.* 2002; 21 (2): 205-222.